

Solving Logarithmic Equations

Name: _____ Date: _____ Score: _____ / 33

Q Quick Review

Solving log equations boils down to one of two moves, plus a mandatory domain check at the end.

Move 1: single log = constant. Get the equation into the form $\log_b(\text{stuff}) = c$, then rewrite in exponential form $\text{stuff} = b^c$ and solve.

Move 2: log = log, same base. If both sides reduce to the form $\log_b(A) = \log_b(B)$, then $A = B$. Logs are one-to-one.

Combining first. When you see multiple log terms, use the product/quotient/power rules to combine them into one log. Once everything is a single log on each side, apply Move 1 or Move 2.

The extraneous-solutions trap — always check. Solving log equations often leads to a polynomial (often a quadratic). Some of its roots may force the original log arguments to be zero or negative, which is undefined for real logs. *Discard those.* Quick check: $\log_2(x) + \log_2(x - 6) = 4$ becomes $x(x - 6) = 16$, so $x^2 - 6x - 16 = 0$, giving $x = 8$ or $x = -2$. Throw out $x = -2$ because the original $\log_2(x)$ needs $x > 0$. The valid answer is $x = 8$.


The check itself is a step — not a verification. Even if the algebra is clean, write down “Check: argument > 0 ? Yes / No” for each original log. If a candidate fails, cross it off; if all fail, the equation has *no solution*.

Common slips. Forgetting the check (most common). Combining logs that don’t share a base. Using $\log(A) + \log(B) = \log(A + B)$ (wrong — it’s $\log(AB)$). Skipping the change-of-base when bases differ on the two sides.

PRACTICE

Combine, exponentiate, solve, then verify the domain. List extraneous solutions explicitly.


1. Solve $\log_3(x) = 4$. _____
2. Solve $\log_2(x + 1) = 5$. _____
3. Solve $\log_2(x) + \log_2(x - 6) = 4$ and discard extraneous roots. _____
4. Solve $\log(x + 5) = \log(2x - 3)$. _____
5. Solve $\log_3(2x + 1) - \log_3(x - 1) = 1$. _____
6. Solve $\log(x) + \log(x - 3) = 1$ and identify any extraneous roots. _____
7. Solve $\ln(x + 4) - \ln(x) = \ln(3)$. _____
8. Which steps below are VALID when solving log equations? Combine same-base logs; convert $\log_b(x) = c$ to $x = b^c$; check domain; equate arguments when $\log_b(A) = \log_b(B)$. _____
9. Solve $\log_5(x) + \log_5(x - 4) = 1$. The number line below shades the valid domain for the original equation — use it to test each candidate root. _____



10. Solve $\log_2(x - 1) + \log_2(x - 5) = 3$. State extraneous solutions. _____
11. Solve $\log(2x) = 3$. _____
12. Solve $\log_4(x) = \frac{3}{2}$. _____
13. Solve $\ln(3x - 2) = \ln(x + 8)$. _____
14. Solve $\log(x^2) = \log(4x - 3)$. _____
15. Solve $\log_2(x) + \log_2(x + 2) = 3$ and discard extraneous roots. _____



16. Solve $\log_3(x + 3) + \log_3(x - 3) = 3$. The valid domain is shown on the number line below; use it to decide which root to keep. _____



17. Solve $\log(x + 2) + \log(x - 1) = \log(4x - 2)$. _____
18. Identify extraneous candidates: $\log(x - 2) + \log(x + 5) = \log(8)$ gives $x = 3$ or $x = -6$. _____
19. Solve $\log_2(8x) = 5$. The table of values for $\log_2(x)$ below may help you read off the step where $\log_2(x) = 2$. _____

x	1	2	8	16	32
$\log_2(x)$	0	1	3	4	5

20. Solve $\log_6(x) + \log_6(x - 5) = 2$. _____

◆ Word Problems

21. Solve $\log_4(x) + \log_4(x - 6) = 2$. Show both roots of the resulting quadratic and explain which one is extraneous and why. _____
22. Solve $\log(x + 3) + \log(x) = \log(28)$. State any extraneous roots explicitly. Then back-substitute to verify. _____
23. Solve $\log_3(x - 2) + \log_3(x + 6) = 2$. List the extraneous root and explain which domain restriction kills it. _____
24. A geology student is told that the equation $\log_2(x + 1) + \log_2(x - 3) = 4$ describes a calibration step in a sensor reading. Solve for x and explain the extraneous candidate in terms of the log's domain. _____

Additional Practice

25. Evaluate $\log_2 32$. _____
26. Evaluate $\log_5 125$. _____
27. Rewrite $\log_3 81 = 4$ exponentially. _____
28. Solve $\log_4 x = 3$. _____
29. Domain of $y = \log(x - 7)$. _____
30. Expand $\log(ab)$. _____
31. Expand $\log(x^3)$. _____
32. Condense $\log x + \log 5$. _____
33. Condense $2 \log x$. _____



Answer Keys

1. $x = 81$
2. $x = 31$
3. $x = 8$ (extraneous: $x = -2$)
4. $x = 8$
5. $x = 4$
6. $x = 5$ (extraneous: $x = -2$)
7. $x = 2$
8. all four
9. $x = 5$ (extraneous: $x = -1$)
10. $x = 3 + 2\sqrt{3}$ (extraneous: $x = 3 - 2\sqrt{3}$)
11. $x = 500$
12. $x = 8$
13. $x = 5$
14. $x = 1$ or $x = 3$
15. $x = 2$ (extraneous: $x = -4$)
16. $x = 6$ (extraneous: $x = -6$)
17. $x = 3$ (extraneous: $x = 0$)
18. $x = 3$ (extraneous: $x = -6$)
19. $x = 4$
20. $x = 9$ (extraneous: $x = -4$)
21. $x = 8$ (extraneous: $x = -2$)
22. $x = 4$ (extraneous: $x = -7$)
23. $x = 3$ (extraneous: $x = -7$)
24. $x = 1 + 2\sqrt{5}$ (extraneous: $x = 1 - 2\sqrt{5}$)

Additional Practice Answers

25. 5
26. 3
27. $3^4 = 81$
28. $x = 64$
29. $x > 7$
30. $\log a + \log b$
31. $3 \log x$
32. $\log(5x)$
33. $\log(x^2)$

Additional Practice: Answers for all numbered items, including the added practice, are shown in the grid above.

Step-by-Step Explanations

1. This is a single log equal to a constant, so rewrite it in exponential form: $x = 3^4 = 81$. Domain check: the argument $x = 81 > 0$, so the solution is valid.
2. Single log equals a constant, so go to exponential form: $x + 1 = 2^5 = 32$. Subtract 1 to get $x = 31$. Domain check: the argument $x + 1 = 32 > 0$, so it is valid.
3. Combine: $\log_2(x(x - 6)) = 4$, so $x(x - 6) = 16$, $x^2 - 6x - 16 = 0$, $(x - 8)(x + 2) = 0$. $x = -2$ makes the original $\log_2(x)$ undefined — toss it. Only $x = 8$ survives.
4. Same base, equate arguments: $x + 5 = 2x - 3$, so $x = 8$. Check both arguments at $x = 8$: $13 > 0$ and $13 > 0 \checkmark$.
5. Quotient: $\log_3\left(\frac{2x+1}{x-1}\right) = 1$. Exponential: $\frac{2x+1}{x-1} = 3$. Cross-multiply: $2x + 1 = 3x - 3$, so $x = 4$. Check: $2x + 1 = 9 > 0$, $x - 1 = 3 > 0 \checkmark$.
6. Combine: $\log(x(x - 3)) = 1$, so $x(x - 3) = 10$, $x^2 - 3x - 10 = 0$, $(x - 5)(x + 2) = 0$. $x = -2$ makes $\log(x)$ undefined; discard. Only $x = 5$ is valid.
7. Quotient: $\ln\left(\frac{x+4}{x}\right) = \ln 3$. Equate arguments: $\frac{x+4}{x} = 3$, so $x + 4 = 3x$ and $x = 2$. Check: $x + 4 = 6 > 0$, $x = 2 > 0 \checkmark$.
8. These are the four legitimate moves. The two illegitimate ones to avoid: $\log A + \log B \neq \log(A + B)$, and $\log A \cdot \log B \neq \log(AB)$.
9. Combine: $\log_5(x(x - 4)) = 1$, so $x(x - 4) = 5$, $x^2 - 4x - 5 = 0$, $(x - 5)(x + 1) = 0$. Original requires $x > 0$ and $x - 4 > 0$, i.e., $x > 4$ — exactly the ray shown. So $x = -1$ fails (left of the open circle); $x = 5$ survives.
10. Combine: $\log_2((x - 1)(x - 5)) = 3$, so $(x - 1)(x - 5) = 8$, $x^2 - 6x - 3 = 0$. Quadratic formula: $x = \frac{6 \pm \sqrt{36 + 12}}{2} = 3 \pm 2\sqrt{3}$. The domain needs $x > 5$. Now $3 - 2\sqrt{3} \approx 0.54$ fails (< 5). $3 + 2\sqrt{3} \approx 6.46$ passes. Only the + root survives.
11. No base is written, so it is base 10. Rewrite in exponential form: $2x = 10^3 = 1000$, then divide by 2 to get $x = 500$. Domain check: the argument $2x = 1000 > 0$, valid.
12. Rewrite in exponential form: $x = 4^{3/2}$. Read the fractional exponent as root-then-power: $4^{3/2} = (\sqrt{4})^3 = 2^3 = 8$. Domain check: $x = 8 > 0$, valid.
13. Equate arguments: $3x - 2 = x + 8$, so $2x = 10$ and $x = 5$. Check: $3(5) - 2 = 13 > 0$ and $5 + 8 = 13 > 0 \checkmark$.

14. Equate arguments: $x^2 = 4x - 3$, so $x^2 - 4x + 3 = 0$ and $(x - 1)(x - 3) = 0$. Check: at $x = 1$, both sides give $\log(1) = 0 \checkmark$. At $x = 3$, both sides give $\log(9) \checkmark$. Both valid.
15. Combine: $\log_2(x(x + 2)) = 3$, so $x(x + 2) = 8$, $x^2 + 2x - 8 = 0$, $(x - 2)(x + 4) = 0$. Original requires $x > 0$, so $x = -4$ is extraneous; $x = 2$ is the answer.
16. Combine: $\log_3(x^2 - 9) = 3$, so $x^2 - 9 = 27$, $x^2 = 36$, $x = \pm 6$. Domain needs $x + 3 > 0$ and $x - 3 > 0$, i.e., $x > 3$ — the ray on the line. So $x = -6$ falls outside it and fails; $x = 6$ wins.
17. Combine LHS: $\log((x + 2)(x - 1)) = \log(4x - 2)$. Equate: $(x + 2)(x - 1) = 4x - 2$, $x^2 + x - 2 = 4x - 2$, $x^2 - 3x = 0$, $x(x - 3) = 0$. So $x = 0$ or $x = 3$. At $x = 0$: $\log(2) + \log(-1)$ — the second log is undefined. At $x = 3$: $\log(5) + \log(2) = \log(10)$ and RHS = $\log(10) \checkmark$. The valid solution is $x = 3$.
18. Keep the rule visible: At $x = -6$: $\log(-8)$ and $\log(-1)$ are both undefined. At $x = 3$: $\log(1) + \log(8) = 0 + \log 8 = \log 8 \checkmark$. That gives a quick check on the answer.
19. One steady path is: $8x = 2^5 = 32$, so $x = 4$. (Or expand first: $3 + \log_2(x) = 5$, so $\log_2(x) = 2$. The table jumps from $\log_2(2) = 1$ to $\log_2(8) = 3$, so $\log_2(x) = 2$ sits between, at $x = 4$. Either way.) That gives a quick check on the answer.
20. Combine: $\log_6(x(x - 5)) = 2$, $x(x - 5) = 36$, $x^2 - 5x - 36 = 0$, $(x - 9)(x + 4) = 0$. Domain: $x > 5$. So $x = -4$ fails, $x = 9$ wins.
21. Combine using the product rule: $\log_4(x(x - 6)) = 2$. Convert to exponential form: $x(x - 6) = 4^2 = 16$. Expand: $x^2 - 6x - 16 = 0$, which factors as $(x - 8)(x + 2) = 0$. So the candidates are $x = 8$ and $x = -2$. Now the domain check: the original equation has $\log_4(x)$ and $\log_4(x - 6)$, so we need both $x > 0$ and $x - 6 > 0$, i.e., $x > 6$. $x = -2$ fails on both counts. $x = 8$ passes ($8 > 6$). So $x = -2$ is extraneous — it would make $\log_4(-2)$ and $\log_4(-8)$, both undefined. The valid solution is $x = 8$. Verify: $\log_4(8) + \log_4(2) = \log_4(16) = 2 \checkmark$.
22. Combine the LHS by the product property: $\log(x(x + 3)) = \log(28)$. Equate arguments: $x(x + 3) = 28$, so $x^2 + 3x - 28 = 0$. Factor: $(x + 7)(x - 4) = 0$. Candidates: $x = -7$ or $x = 4$. Domain check on the original: $\log(x + 3)$ needs $x > -3$, and $\log(x)$ needs $x > 0$, so altogether $x > 0$. $x = -7$ fails ($-7 < 0$, would give $\log(-7)$ and $\log(-4)$ — both undefined). $x = 4$ passes. Back-substitute $x = 4$: $\log(7) + \log(4) = \log(28) \checkmark$ by the product rule. (Always do this final plug-back step. It catches arithmetic slips, too.)
23. Combine: $\log_3((x - 2)(x + 6)) = 2$. Exponential form: $(x - 2)(x + 6) = 9$.



Scan Me

Expand: $x^2 + 4x - 12 = 9$, so $x^2 + 4x - 21 = 0$ and $(x + 7)(x - 3) = 0$.
 Candidates: $x = -7$ or $x = 3$. Domain: $x - 2 > 0$ requires $x > 2$, and $x + 6 > 0$
 requires $x > -6$. Both must hold, so $x > 2$. $x = -7$ fails the stricter condition
 $x > 2$ (and also makes $x - 2 = -9$, i.e., $\log_3(-9)$ undefined). $x = 3$ passes
 ($3 > 2$). Verify: $\log_3(1) + \log_3(9) = 0 + 2 = 2 \checkmark$.


24. Combine: $\log_2((x+1)(x-3)) = 4$. Exponential form: $(x+1)(x-3) = 2^4 = 16$.
 Expand: $x^2 - 2x - 3 = 16$, so $x^2 - 2x - 19 = 0$. Use the quadratic formula:

$x = \frac{2 \pm \sqrt{4 + 76}}{2} = \frac{2 \pm \sqrt{80}}{2} = 1 \pm 2\sqrt{5}$. Numerically: $1 + 2\sqrt{5} \approx 5.47$
 and $1 - 2\sqrt{5} \approx -3.47$. Domain: $x + 1 > 0$ and $x - 3 > 0$, so $x > 3$. The
 positive root ≈ 5.47 passes; the negative root fails. So $x = 1 + 2\sqrt{5} \approx 5.47$
 is the valid solution; $1 - 2\sqrt{5} \approx -3.47$ is extraneous. (The reading $x \approx 5.5$ is
 the physically meaningful one; the negative candidate is an arithmetic shadow the
 algebra produces but the log forbids.)



Keep Building ALEKS Skills

Recommended ALEKS books and bundle




The Most Comprehensive ALEKS Math Preparation Bundle

Prep books, workbooks, full-length tests, detailed explanations, and a formula bonus file in one complete ALEKS study path.



Scan Me
Download Instantly

STUDENT FAVORITE · ALEKS Math for Beginners



ALEKS Math for Beginners 2026

Step-by-step lessons, basic study guides, topic practice, and two full-length ALEKS Math tests for students who need a calm path from review to readiness.

Great companion to these worksheets for self-study, tutoring, and classroom review.

PDF Edition



Scan Me

Amazon



Scan Me

For more ALEKS prep, visit EffortlessMath.com/ALEKS