

The Discriminant

Name: _____ Date: _____ Score: _____ / 24

Quick Review

For a quadratic equation $ax^2 + bx + c = 0$, the **discriminant** is $D = b^2 - 4ac$. It tells you the number and type of solutions before you solve. If $D > 0$, there are two real solutions; if $D = 0$, there is one real solution; if $D < 0$, there are no real solutions. If D is a perfect square, the real solutions are rational; if D is positive but not a perfect square, the real solutions are irrational.

PRACTICE

Find the discriminant and describe the solutions.

- | | | | |
|-------------------------|-------|--------------------------------------|-------|
| 1. $x^2 - 5x + 6 = 0$ | _____ | 11. $6x^2 + x + 2 = 0$ | _____ |
| 2. $x^2 + 4x + 4 = 0$ | _____ | 12. $9x^2 - 12x + 4 = 0$ | _____ |
| 3. $x^2 + x + 1 = 0$ | _____ | 13. $x^2 + 6x + 5 = 0$ | _____ |
| 4. $2x^2 - 3x - 2 = 0$ | _____ | 14. $x^2 + 2x - 3 = 0$ | _____ |
| 5. $3x^2 + 2x + 5 = 0$ | _____ | 15. $2x^2 + 4x + 7 = 0$ | _____ |
| 6. $x^2 - 2x - 1 = 0$ | _____ | 16. $3x^2 - 10x + 2 = 0$ | _____ |
| 7. $4x^2 + 4x + 1 = 0$ | _____ | 17. $x^2 - 8x + 16 = 0$ | _____ |
| 8. $5x^2 - x + 2 = 0$ | _____ | 18. $7x^2 + 2x - 1 = 0$ | _____ |
| 9. $x^2 - 9 = 0$ | _____ | 19. $x^2 + 5x + 8 = 0$ | _____ |
| 10. $2x^2 + 7x + 3 = 0$ | _____ | 20. If $D = 49$, describe solutions | _____ |

Word Problems

21. A quadratic model has equation $h(t) = -16t^2 + 32t + 4$. Use the discriminant to decide whether the object reaches ground level.

22. A profit equation is $P(x) = -x^2 + 6x - 12$. Use the discriminant to decide whether profit ever equals zero.

23. A parabola touches the x -axis at exactly one point. What must be true about its discriminant?

24. A quadratic equation has discriminant 18. How many real solutions does it have, and are they rational?



Answer Keys

- | | |
|--|---|
| <p>1. $D = 1$; two rational</p> <p>2. $D = 0$; one real</p> <p>3. $D = -3$; no real</p> <p>4. $D = 25$; two rational</p> <p>5. $D = -56$; no real</p> <p>6. $D = 8$; two irrational</p> <p>7. $D = 0$; one real</p> <p>8. $D = -39$; no real</p> <p>9. $D = 36$; two rational</p> <p>10. $D = 25$; two rational</p> <p>11. $D = -47$; no real</p> <p>12. $D = 0$; one real</p> | <p>13. $D = 16$; two rational</p> <p>14. $D = 16$; two rational</p> <p>15. $D = -40$; no real</p> <p>16. $D = 76$; two irrational</p> <p>17. $D = 0$; one real</p> <p>18. $D = 32$; two irrational</p> <p>19. $D = -7$; no real</p> <p>20. two rational</p> <p>21. $D = 1280$; yes, two real times</p> <p>22. $D = -12$; no real break-even point</p> <p>23. $D = 0$</p> <p>24. two real irrational solutions</p> |
|--|---|


Step-by-Step Tutor Notes

1. This is a good place to slow down, check the notation, and simplify cleanly. $D = (-5)^2 - 4(1)(6) = 25 - 24 = 1$. So the answer is $D = 1$; two rational.
2. Start with the definition the problem is testing, then apply it directly. $D = 16 - 16 = 0$. So the answer is $D = 0$; one real.
3. Start with the definition the problem is testing, then apply it directly. $D = 1 - 4 = -3$. So the answer is $D = -3$; no real.
4. Start with the definition the problem is testing, then apply it directly. $D = (-3)^2 - 4(2)(-2) = 9 + 16 = 25$. So the answer is $D = 25$; two rational.
5. Start with the definition the problem is testing, then apply it directly. $D = 4 - 60 = -56$. So the answer is $D = -56$; no real.
6. This is a good place to slow down, check the notation, and simplify cleanly. $D = 4 + 4 = 8$, positive but not a perfect square. So the answer is $D = 8$; two irrational.
7. Start with the definition the problem is testing, then apply it directly. $D = 16 - 16 = 0$. So the answer is $D = 0$; one real.
8. Focus on the main idea of the problem, then simplify carefully. $D = 1 - 40 = -39$. So the answer is $D = -39$; no real.
9. This is a good place to slow down, check the notation, and simplify cleanly. Here $a = 1, b = 0, c = -9$, so $D = 36$. So the answer is $D = 36$; two rational.
10. Focus on the main idea of the problem, then simplify carefully. $D = 49 - 24 = 25$. So the answer is $D = 25$; two rational.
11. Use the clue in the question first, then let the arithmetic finish the job. $D = 1 - 48 = -47$. So the answer is $D = -47$; no real.
12. Start with the definition the problem is testing, then apply it directly. $D = 144 - 144 = 0$. So the answer is $D = 0$; one real.
13. This is a good place to slow down, check the notation, and simplify cleanly. $D = 36 - 20 = 16$. So the answer is $D = 16$; two rational.
14. Start with the definition the problem is testing, then apply it directly. $D = 4 + 12 = 16$. So the answer is $D = 16$; two rational.
15. Focus on the main idea of the problem, then simplify carefully. $D = 16 - 56 = -40$. So the answer is $D = -40$; no real.
16. This is a good place to slow down, check the notation, and simplify cleanly. $D = 100 - 24 = 76$. So the answer is $D = 76$; two irrational.
17. Take it one clear step at a time and keep the original question in mind. $D = 64 - 64 = 0$. So the answer is $D = 0$; one real.
18. Focus on the main idea of the problem, then simplify carefully. $D = 4 + 28 = 32$. So the answer is $D = 32$; two irrational.
19. Take it one clear step at a time and keep the original question in mind. $D = 25 - 32 = -7$. So the answer is $D = -7$; no real.
20. Start with the definition the problem is testing, then apply it directly. Positive perfect square means two rational solutions. So the answer is two rational.
21. Set $h = 0$. With $a = -16, b = 32, c = 4, D = 32^2 - 4(-16)(4) = 1280 > 0$, so the graph crosses the ground level.
22. Here $a = -1, b = 6, c = -12$. The discriminant is $36 - 48 = -12$, so the model has no real zeros.
23. Touching once means the quadratic has one real solution, which happens exactly when the discriminant is 0.
24. 18 is positive, so there are two real solutions. It is not a perfect square, so the solutions are irrational.



Keep Building ALEKS Skills

Recommended ALEKS books and bundle




The Most Comprehensive ALEKS Math Preparation Bundle

Prep books, workbooks, full-length tests, detailed explanations, and a formula bonus file in one complete ALEKS study path.



Scan Me
Download Instantly

STUDENT FAVORITE · ALEKS Math for Beginners



ALEKS Math for Beginners 2026

Step-by-step lessons, basic study guides, topic practice, and two full-length ALEKS Math tests for students who need a calm path from review to readiness.

Great companion to these worksheets for self-study, tutoring, and classroom review.

PDF Edition



Scan Me

Amazon



Scan Me

For more ALEKS prep, visit EffortlessMath.com/ALEKS