

# Rational Expressions

Name: \_\_\_\_\_ Date: \_\_\_\_\_ Score: \_\_\_\_\_ / 18

## Quick Review and Helpful Hints

A *rational expression* is a fraction whose top and bottom are polynomials. Simplify by factoring both parts and canceling common factors. It is *undefined* wherever the denominator equals 0, so those *x*-values must be excluded.

▶ **Example:** Simplify  $\frac{x^2 - 9}{x + 3}$ . **Work:** Factor the top:  $\frac{(x + 3)(x - 3)}{x + 3}$ .  
 Cancel the common  $(x + 3)$ .      ★ **Answer:**  $x - 3$  ( $x \neq -3$ )

$$\frac{\cancel{(x + 3)}(x - 3)}{\cancel{(x + 3)}} = x - 3$$

Cancel common factors.

### ◆ Practice Problems

Simplify each expression (or answer as directed).

- |   |  |
|---|--|
| <p>1. <math>\frac{2x}{4}</math> _____</p> <p>2. <math>\frac{6x^2}{3x}</math> _____</p> <p>3. <math>\frac{x^2 - 4}{x - 2}</math> _____</p> <p>4. <math>\frac{x^2 - 9}{x + 3}</math> _____</p> <p>5. <math>\frac{x^2 + 5x + 6}{x + 2}</math> _____</p> <p>6. <math>\frac{x^2 - x - 6}{x - 3}</math> _____</p> <p>7. <math>\frac{3x + 6}{3}</math> _____</p> | <p>8. <math>\frac{x^2 - 16}{x - 4}</math> _____</p> <p>9. <math>\frac{5x^2}{10x}</math> _____</p> <p>10. <math>\frac{x^2 + 7x + 10}{x + 5}</math> _____</p> <p>11. <math>\frac{x^2 - 25}{x + 5}</math> _____</p> <p>12. <math>\frac{4x + 8}{x + 2}</math> _____</p> <p>13. <math>\frac{x^2 - 1}{x - 1}</math> _____</p> <p>14. <math>\frac{1}{x - 3}</math> is undefined when <math>x =</math> _____</p> |
|---|--|

### ◆ Word Problems

- |  |  |
|--|--|
| <p>15. A rectangle has area <math>x^2 - 36</math> and one side length <math>x + 6</math>. Simplify <math>\frac{x^2 - 36}{x + 6}</math> to represent the other side length. _____</p> <p>16. Where is <math>\frac{5}{x - 2}</math> undefined? _____</p> | <p>17. A batch recipe has total ingredient amount <math>2x^2 + 4x</math> split evenly across <math>2x</math> containers. Simplify <math>\frac{2x^2 + 4x}{2x}</math> for the amount per container. _____</p> <p>18. A rectangular garden has area <math>x^2 + 8x + 15</math> and one side <math>x + 3</math>. Simplify <math>\frac{x^2 + 8x + 15}{x + 3}</math> for the missing side. _____</p> |
|--|--|



## Answer Keys

1.  $\frac{x}{2}$

2.  $2x$

3.  $x + 2$

4.  $x - 3$

5.  $x + 3$

6.  $x + 2$

7.  $x + 2$

8.  $x + 4$

9.  $\frac{x}{2}$

10.  $x + 2$

11.  $x - 5$

12.  $4$

13.  $x + 1$

14.  $3$

15.  $x - 6$

16.  $x = 2$

17.  $x + 2$

18.  $x + 5$

### Step-by-Step Explanations

1. Start by naming the process: For a rational expression, factor first when possible, then cancel only common factors from the top and bottom. The setup/work is Divide top and bottom by 2:  $\frac{x}{2}$ . So the final answer is  $\frac{x}{2}$ .

2. A good way to think about this is: For a rational expression, factor first when possible, then cancel only common factors from the top and bottom. The setup/work is  $\frac{6x^2}{3x} = 2x$ . So the final answer is  $2x$ .

3. Step by step: For a rational expression, factor first when possible, then cancel only common factors from the top and bottom. The setup/work is  $\frac{(x-2)(x+2)}{x-2} = x+2$ . So the final answer is  $x+2$ .

4. Take it one move at a time: For a rational expression, factor first when possible, then cancel only common factors from the top and bottom. The setup/work is  $\frac{(x+3)(x-3)}{x+3} = x-3$ . So the final answer is  $x-3$ .

5. Start by naming the process: For a rational expression, factor first when possible, then cancel only common factors from the top and bottom. The setup/work is  $\frac{(x+2)(x+3)}{x+2} = x+3$ . So the final answer is  $x+3$ .

6. A good way to think about this is: For a rational expression, factor first when possible, then cancel only common factors from the top and bottom. The setup/work is  $\frac{(x-3)(x+2)}{x-3} = x+2$ . So the final answer is  $x+2$ .

7. Step by step: For a rational expression, factor first when possible, then cancel only common factors from the top and bottom. The setup/work is  $\frac{3(x+2)}{3} = x+2$ . So the final answer is  $x+2$ .

8. Take it one move at a time: For a rational expression, factor first when possible, then cancel only common factors from the top and bottom. The setup/work is  $\frac{(x-4)(x+4)}{x-4} = x+4$ . So the final answer is  $x+4$ .

9. Start by naming the process: For a rational expression, factor first when possible, then cancel only common factors from the top and bottom. The setup/work is  $\frac{5x^2}{10x} = \frac{x}{2}$ . So the final answer is  $\frac{x}{2}$ .

10. A good way to think about this is: For a rational expression, factor first when possible, then cancel only common factors from the top and bottom. The setup/work is  $\frac{(x+5)(x+2)}{x+5} = x+2$ . So the final answer is  $x+2$ .

11. Step by step: For a rational expression, factor first when possible, then cancel only common factors from the top and bottom. The setup/work is  $\frac{(x-5)(x+5)}{x+5} = x-5$ . So the final answer is  $x-5$ .

12. Take it one move at a time: For a rational expression, factor first when possible, then cancel only common factors from the top and bottom. The setup/work is  $\frac{4(x+2)}{x+2} = 4$ . So the final answer is 4.

13. Start by naming the process: For a rational expression, factor first when possible, then cancel only common factors from the top and bottom. The setup/work is  $\frac{(x-1)(x+1)}{x-1} = x+1$ . So the final answer is  $x+1$ .

14. A good way to think about this is: For a rational expression, factor first when possible, then cancel only common factors from the top and bottom. The setup/work is The bottom is 0 when  $x = 3$ . So the final answer is 3.

15. Step by step: For a rational expression, factor first when possible, then cancel only common factors from the top and bottom. The setup/work is  $\frac{(x-6)(x+6)}{x+6} = x-6$ . So the final answer is  $x-6$ .

16. Take it one move at a time: For a rational expression, factor first when possible, then cancel only common factors from the top and bottom. The setup/work is The denominator is 0 at  $x = 2$ . So the final answer is  $x = 2$ .


17. Start by naming the process: For a rational expression, factor first when possible, then cancel only common factors from the top and bottom. The setup/work is  $\frac{2x(x+2)}{2x} = x+2$ . So the final answer is  $x+2$ .

18. A good way to think about this is: For a rational expression, factor first when possible, then cancel only common factors from the top and bottom. The setup/work is  $\frac{(x+3)(x+5)}{x+3} = x+5$ . So the final answer is  $x+5$ .



# Keep Building ASVAB Math Skills

Recommended Effortless Math resources




## The Most Comprehensive ASVAB Math Preparation Bundle

Use the complete ASVAB Math resource for review, worked examples, extra practice, and test-style questions after each worksheet.



Scan Me  
Download Instantly

## STUDENT FAVORITE - ASVAB Math for Beginners



## ASVAB Math for Beginners 2026

Step-by-step lessons, topic practice, and full review support for students who want a calm path through ASVAB Math preparation.

A strong companion for self-study, tutoring, homework, and targeted review.

PDF Edition



Scan Me  
Download Instantly

For more ASVAB Math prep, visit [EffortlessMath.com/ASVAB](https://www.EffortlessMath.com/ASVAB)