

Adding and Subtracting Functions

Name: _____ Date: _____ Score: _____ / 37

Q Quick Review

You can **add** or **subtract** two functions by working on their outputs at each input. The definitions: $(f + g)(x) = f(x) + g(x)$ and $(f - g)(x) = f(x) - g(x)$. The new function gets a new name (often $f + g$ or $f - g$), but the rule is just “combine the outputs.”

Watch the minus. When subtracting $(f - g)(x)$, distribute the negative across *every term* of g . So if $f(x) = 2x^2 - 3x + 5$ and $g(x) = x^2 + 4x - 1$, then $(f - g)(x) = (2x^2 - 3x + 5) - (x^2 + 4x - 1) = 2x^2 - 3x + 5 - x^2 - 4x + 1 = x^2 - 7x + 6$. The $+1$ at the end is the sign-flip many students miss.

Domains. The domain of $f + g$ (or $f - g$) is the *intersection* of the domains of f and g . If $f(x) = \sqrt{x-1}$ (needs $x \geq 1$) and $g(x) = \sqrt{4-x}$ (needs $x \leq 4$), then $(f + g)(x)$ is defined only on $[1, 4]$.

Properties. Addition is commutative: $f + g = g + f$. Subtraction isn't — $f - g$ and $g - f$ are negatives of each other. To evaluate at a number, you can build the combined formula first or just compute $f(c)$ and $g(c)$ separately and combine. The second route is usually faster.

PRACTICE

Combine the functions as indicated. Simplify and state any domain restrictions when asked.

1. $f(x) = 2x + 3, g(x) = x - 1; (f + g)(x)$ _____
2. $f(x) = 4x + 5, g(x) = x + 2; (f - g)(x)$ _____
3. $f(x) = x^2 + 3x - 1, g(x) = 2x + 4; (f + g)(x)$ _____
4. $f(x) = 2x^2 - 3x + 5, g(x) = x^2 + 4x - 1; (f - g)(x)$ _____
5. The tables give f and g . Find $(f + g)(2)$. _____

x	0	1	2	3
$f(x)$	1	2	5	10

x	0	1	2	3
$g(x)$	-2	1	4	7

6. $f(x) = 3x - 5, g(x) = x^2 + 2x; (f + g)(x)$ _____
7. $f(x) = 3x - 5, g(x) = x^2 + 2x; (f - g)(x)$ _____
8. $f(x) = 5x^2 - 2x + 7, g(x) = 3x^2 + x - 4; (f - g)(x)$ _____
9. $f(x) = \sqrt{x-1}, g(x) = \sqrt{4-x}$; domain of $(f + g)(x)$ _____
10. $f(x) = \sqrt{x+2}, g(x) = \sqrt{6-x}$; domain of $(f + g)(x)$ _____
11. $f(x) = x^3 + 2x, g(x) = x^3 - x; (f - g)(x)$ _____
12. $f(x) = 4x^2 + x, g(x) = 4x^2 - x; (f + g)(x)$ _____
13. $f(x) = 2x + 1, g(x) = 3x - 4; (f - g)(5)$ _____
14. $f(x) = x^2, g(x) = 2x + 3$; find $(f - g)(-1)$ _____



15. The tables give f and g . Find $(f + g)(0)$. _____

x	-1	0	1	2
$f(x)$	-1	2	5	8

x	-1	0	1	2
$g(x)$	0	-1	0	3

16. $R(x) = 45x + 200$ revenue, $C(x) = 18x + 125$ cost. Profit $P = R - C$. _____

17. $f(x) = 2x - 7$, $g(x) = 5 - x$; find x with $(f + g)(x) = 0$ _____

18. $A + B = 5x^2 + 3x - 2$, $A = 2x^2 - x + 4$. Find B . _____

19. The tables give f and g . Find $(f - g)(3)$. _____

x	1	2	3	4
$f(x)$	-3	0	5	12

x	1	2	3	4
$g(x)$	3	4	5	6

20. $f(x) = |x|$, $g(x) = x$. For $x < 0$, simplify $(f - g)(x)$. _____

◆ Word Problems

21. A store's revenue is modeled by $R(x) = 45x + 200$ dollars and its cost is $C(x) = 18x + 125$ dollars, where x is the number of items sold. Write the profit function $P(x) = R(x) - C(x)$ and find $P(10)$. _____

22. Two divisions of a company contribute to quarterly profit. Division X contributes $X(t) = 2t^2 - 3t + 5$ thousand dollars and division Y contributes $Y(t) = -t^2 + 4t + 3$ thousand dollars, where t is months since the quarter started. Write the total profit function $T(t)$ in simplified form. _____

23. Let $f(x) = \sqrt{x+3}$ and $g(x) = \sqrt{5-x}$. Find the domain of $(f - g)(x)$, and evaluate it at $x = 1$. _____

24. A garden's total perimeter is $T(x) = 8x + 14$ meters, made up of two pieces: the front fence $F(x) = 3x + 4$ meters and the side fences combined. Write a function $S(x)$ for the total side-fence length, then find $S(5)$. _____

Additional Practice

25. If $f(x) = 2x - 5$, find $f(4)$. _____

26. If $g(x) = x^2 + 1$, find $g(-3)$. _____

27. For $f(x) = 3x + 2$, solve $f(x) = 14$. _____

28. Find $(f + g)(x)$ if $f = x + 1$, $g = 2x - 5$. _____

29. Find $(fg)(x)$ if $f = x - 2$, $g = x + 3$. _____

30. Find $f(g(x))$ if $f(x) = 2x$, $g(x) = x + 7$. _____

31. Find the inverse of $f(x) = x - 9$. _____

32. Find the inverse of $f(x) = 3x + 1$. _____



33. Domain of $f(x) = \sqrt{x - 4}$. _____

34. Domain of $f(x) = \frac{1}{x + 6}$. _____

35. Parent function for $y = |x| + 3$. _____

36. Shift $y = x^2$ left 4. _____

37. Average rate from $(1, 5)$ to $(4, 17)$. _____



Answer Keys

1. $3x + 2$
 2. $3x + 3$
 3. $x^2 + 5x + 3$
 4. $x^2 - 7x + 6$
 5. 9
 6. $x^2 + 5x - 5$
 7. $-x^2 + x - 5$
 8. $2x^2 - 3x + 11$
 9. [1, 4]
 10. [-2, 6]
 11. $3x$
 12. $8x^2$
 13. 0
 14. 0
 15. 1
 16. $P(x) = 27x + 75$
 17. $x = 2$
 18. $B = 3x^2 + 4x - 6$
 19. 0
 20. $-2x$
 21. $P(x) = 27x + 75$, $P(10) = \$345$
 22. $T(t) = t^2 + t + 8$ thousand dollars
 23. domain [-3, 5]; $(f - g)(1) = 0$
 24. $S(x) = 5x + 10$, $S(5) = 35$ meters

Additional Practice Answers

25. 3
 26. 10
 27. $x = 4$
 28. $3x - 4$
 29. $x^2 + x - 6$
 30. $2x + 14$
 31. $f^{-1}(x) = x + 9$
 32. $f^{-1}(x) = \frac{x-1}{3}$
 33. $x \geq 4$
 34. $x \neq -6$
 35. $y = |x|$
 36. $y = (x + 4)^2$
 37. 4

Additional Practice: Answers for all numbered items, including the added practice, are shown in the grid above.

Step-by-Step Explanations

1. Add the outputs: $(f + g)(x) = (2x + 3) + (x - 1)$. Group like terms — the x -terms give $2x + x = 3x$, the constants give $3 - 1 = 2$. So $(f + g)(x) = 3x + 2$.
 2. Keep the rule visible: Distribute the minus: $(4x + 5) - (x + 2) = 4x + 5 - x - 2 = 3x + 3$. This is the part to check before moving on, because it keeps the answer tied to the original question.
 3. Add the outputs and group like terms: $(x^2 + 3x - 1) + (2x + 4)$. The x^2 has no partner, the x -terms give $3x + 2x = 5x$, and the constants give $-1 + 4 = 3$. So $(f + g)(x) = x^2 + 5x + 3$.
 4. Flip every sign in g : $-(x^2 + 4x - 1) = -x^2 - 4x + 1$. Combine: $(2 - 1)x^2 + (-3 - 4)x + (5 + 1) = x^2 - 7x + 6$. The $+1$ at the end is the easy slip.
 5. Read each table at $x = 2$: $f(2) = 5$ and $g(2) = 4$. Add the outputs: $(f + g)(2) = 5 + 4 = 9$. Reading values straight off a table beats rebuilding the combined formula.
 6. Add the outputs: $(3x - 5) + (x^2 + 2x)$. The x^2 stands alone, the x -terms give $3x + 2x = 5x$, and the only constant is -5 . So $(f + g)(x) = x^2 + 5x - 5$.
 7. One steady path is: Flip g 's signs: $-(x^2 + 2x) = -x^2 - 2x$. Combine: $-x^2 + (3 - 2)x - 5 = -x^2 + x - 5$. That gives a quick check on the answer.
 8. Subtract by flipping every sign in g : $-(3x^2 + x - 4) = -3x^2 - x + 4$. Now combine with f : $(5 - 3)x^2 + (-2 - 1)x + (7 + 4) = 2x^2 - 3x + 11$. The trap is the last term — $-(-4)$ becomes $+4$, so $7 + 4 = 11$, not 3.
 9. For the sum to exist, both square roots must be defined at once. From $\sqrt{x - 1}$ we need $x - 1 \geq 0$, so $x \geq 1$; from $\sqrt{4 - x}$ we need $4 - x \geq 0$, so $x \leq 4$. The domain is where both hold — the intersection [1, 4].
 10. Both radicals must be nonnegative simultaneously. From $\sqrt{x + 2}$: $x + 2 \geq 0$, so $x \geq -2$. From $\sqrt{6 - x}$: $6 - x \geq 0$, so $x \leq 6$. The domain is the overlap of these, the interval [-2, 6].
 11. Subtract by flipping g 's signs: $-(x^3 - x) = -x^3 + x$. Combine with f : the cubes give $x^3 - x^3 = 0$, and the linear terms give $2x + x = 3x$. So $(f - g)(x) = 3x$ — the cubes cancel cleanly.
 12. Add the outputs: $(4x^2 + x) + (4x^2 - x)$. The squared terms give $4x^2 + 4x^2 = 8x^2$, and the linear terms cancel: $x + (-x) = 0$. So $(f + g)(x) = 8x^2$.

13. A careful way to see it: $f(5) = 11$, $g(5) = 11$. Difference: 0. The two functions meet at $x = 5$, so the subtraction gives zero there. That gives a quick check on the answer.
 14. Keep the rule visible: $f(-1) = 1$, $g(-1) = 1$. Difference: $1 - 1 = 0$. (The parabola and the line cross at $x = -1$.) That gives a quick check on the answer.
 15. Look down the $x = 0$ column in each table: $f(0) = 2$ and $g(0) = -1$. Sum the outputs: $(f + g)(0) = 2 + (-1) = 1$.
 16. Distribute the minus: $(45x + 200) - (18x + 125) = 45x + 200 - 18x - 125 = 27x + 75$. A student who adds instead of subtracts gets $63x + 325$ — wrong direction.
 17. A careful way to see it: $(f + g)(x) = (2x - 7) + (5 - x) = x - 2$. Set $x - 2 = 0$: $x = 2$. This is the part to check before moving on, because it keeps the answer tied to the original question.
 18. Keep the rule visible: $B = (A + B) - A = (5x^2 + 3x - 2) - (2x^2 - x + 4)$. Flip A 's signs: $-2x^2 + x - 4$. Combine: $3x^2 + 4x - 6$. That gives a quick check on the answer.
 19. Read at $x = 3$: $f(3) = 5$ and $g(3) = 5$. Subtract: $(f - g)(3) = 5 - 5 = 0$. (The two functions meet at $x = 3$, so their difference is zero there.)
 20. For $x < 0$, $|x| = -x$. So $(f - g)(x) = -x - x = -2x$, which is positive when $x < 0$. (At $x = -3$, $(f - g)(-3) = 3 - (-3) = 6 = -2(-3)$. Sanity-check passes.)
 21. Subtract carefully: $P(x) = (45x + 200) - (18x + 125) = 27x + 75$. Each item sold adds \$27 to profit (the \$45 revenue minus the \$18 cost per item), starting from a \$75 baseline. At $x = 10$: $P(10) = 270 + 75 = 345$. So selling 10 items yields \$345 profit.
 22. Add term by term: $X(t) + Y(t) = (2 - 1)t^2 + (-3 + 4)t + (5 + 3) = t^2 + t + 8$. At the start of the quarter ($t = 0$), total profit is \$8,000 — a sensible baseline. The t^2 term means profit grows faster as the months tick by.
 23. Both radicands must be nonnegative. From f : $x + 3 \geq 0$ so $x \geq -3$. From g : $5 - x \geq 0$ so $x \leq 5$. The intersection is [-3, 5]. At $x = 1$: $f(1) = \sqrt{4} = 2$ and $g(1) = \sqrt{4} = 2$, so $(f - g)(1) = 0$. (The two functions cross at $x = 1$ — their graphs meet there.)
 24. Side fences total $S = T - F = (8x + 14) - (3x + 4)$. Flip the signs in F : $-3x - 4$. Combine: $5x + 10$. At $x = 5$: $S(5) = 25 + 10 = 35$ meters. Sanity-check: $F(5) + S(5) = 19 + 35 = 54 = T(5)$. Numbers add up.



Build Algebra Confidence From Pre-Algebra Through Algebra II



The Complete Algebra Success Bundle

Pre-Algebra, Algebra I, and Algebra II in one clear path

Friendly lessons, focused practice, and full-review support for every stage.



Scan for the Bundle

6 Books
3 Courses
1 Path

Bundle Value: Six coordinated books help students review missing skills, learn new algebra topics, and practice until the steps feel natural.

Complete Course Path

- ✓ Starts with Pre-Algebra foundations
- ✓ Moves smoothly into Algebra I skills
- ✓ Extends learning through Algebra II topics
- ✓ Great for review, tutoring, and summer study

One bundle, one steady path.

Step-by-Step Lessons

- ✓ Plain-English explanations students can follow
- ✓ Worked examples that show every important step
- ✓ Common mistakes called out before they stick
- ✓ Skill-building practice after each lesson
- ✓ Helpful for independent study or class support

Less guessing. More understanding.

Practice That Sticks

- ✓ Matching practice workbooks for extra repetition
- ✓ Review sets to keep older skills fresh
- ✓ Answer explanations for checking thinking
- ✓ Strong support before tests and final exams
- ✓ Designed to build fluency and confidence

Practice today. Remember tomorrow.

STUDENT FAVORITE • Master Algebra II From the Ground Up



Algebra II for Beginners

Written by a top math teacher & aligned with national and state Algebra II courses. From polynomial functions to logarithms, trigonometry, and rational expressions — explained the easy way.

- ✓ **Complete coverage** of every Algebra II concept — perfect companion to these worksheets
- ✓ **Step-by-step explanations** with worked examples on every topic
- ✓ **QR codes in every chapter** for free video lessons & bonus practice
- ✓ **2 full-length practice tests** with detailed answer keys

- ✓ 100% Guaranteed
- ✓ Lifetime Support
- ✓ Trusted by Teachers

Start Your Algebra Journey Today! →

★ STUDENT'S #1 CHOICE ★

Teacher-recommended • 12,000+ Happy Students

PDF EDITION



Instant download • any device

PAPERBACK



Paperback on Amazon

Hold it in your hands

Pair these free worksheets with *Algebra II for Beginners* and you have a complete self-paced course — concept lessons, daily practice, and full exam-style reviews, all in one path. → EffortlessMath.com/product/algebra-ii-for-beginners