

Adding and Subtracting Rational Expressions

Name: _____ Date: _____ Score: _____ / 34

Q Quick Review

Adding or subtracting rational expressions is the same logic as adding numeric fractions: you need a **common denominator** first. With polynomials the LCD takes one extra step (factor each denominator), but the rule is identical.

Step 1 – factor every denominator. Once you see the factors, the LCD is the product of each distinct factor at the highest power it appears in any denominator. Quick check: denominators $x - 3$ and $x^2 - 9 = (x - 3)(x + 3)$ share $(x - 3)$, so the LCD is $(x - 3)(x + 3)$.

Step 2 – rewrite each fraction with the LCD. Multiply numerator and denominator of each piece by whatever factor is missing. Keep the new numerators in factored form – you’ll need to combine them next.

Step 3 – combine the numerators. Add or subtract across the top. The *biggest* trap on subtraction: the negative sign hits *every term* in the second numerator, not just the first. Use parentheses and distribute carefully.

Step 4 – simplify and state restrictions. Factor the resulting numerator if you can, look for any common factor with the LCD, and cancel. State $x \neq$ values from *every* original denominator (whether or not those factors cancel).

What you should never do. Add numerators when denominators differ. That’s the equivalent of saying $\frac{1}{2} + \frac{1}{3} = \frac{2}{5}$ – the universal warning sign of fraction trouble.

The subtraction sign trap. $\frac{a}{d} - \frac{b+c}{d} = \frac{a-b-c}{d}$, not $\frac{a-b+c}{d}$. The minus distributes through the entire second numerator. This catches more students than any other sub-step in the topic.

PRACTICE

Factor denominators, build the LCD, rewrite with it, combine numerators, then simplify and state restrictions.

1. Add $\frac{2}{x} + \frac{3}{x}$. The table evaluates the sum at a few inputs. _____

x	1	2	5
$\frac{2}{x} + \frac{3}{x}$	5	$\frac{5}{2}$	1

2. $\frac{x+5}{x-1} - \frac{x+2}{x-1}$ _____

3. Add $\frac{1}{x} + \frac{2}{x+1}$. The table samples the sum away from the excluded values. _____

x	1	2	3
$\frac{1}{x} + \frac{2}{x+1}$	2	$\frac{7}{6}$	$\frac{5}{6}$

4. Subtract $\frac{3}{x+2} - \frac{2}{x-1}$. The table evaluates the difference at several inputs. _____

x	0	2	3
difference	$\frac{7}{2}$	$-\frac{5}{4}$	$-\frac{2}{5}$

5. What is the LCD of $\frac{1}{x-3}$ and $\frac{2}{x^2-9}$? _____

6. $\frac{1}{x} + \frac{1}{x+1} - \frac{1}{x-1}$ _____

7. $\frac{2}{x-3} + \frac{x}{x+3}$ _____

8. $\frac{3}{x-4} + \frac{2}{x^2-16}$ _____

9. $\frac{x}{x-2} - \frac{5}{x+1}$ _____



10. $\frac{3}{x+1} - \frac{1}{x}$ _____

11. $\frac{5}{x^2-1} + \frac{3}{x+1}$ _____

12. $\frac{2x}{x-1} + \frac{x+1}{x-1}$ _____

13. $\frac{1}{x+2} + \frac{1}{x-2}$ _____

14. $\frac{x+3}{x-2} - \frac{x-2}{x+3}$ _____

15. $\frac{4}{x} - \frac{1}{2x}$ _____

16. Mark TRUE or FALSE: $\frac{a}{b} + \frac{c}{d} = \frac{a+c}{b+d}$. _____

17. $\frac{2x}{x+3} - \frac{x-1}{x+3}$ _____

18. $\frac{x}{x^2-4} + \frac{1}{x-2}$ _____

19. $\frac{5}{x-3} + \frac{4}{3-x}$ _____

20. $\frac{x+1}{x} + \frac{1}{x+1}$ _____

◆ Word Problems

21. Two pipes fill a tank. Pipe A fills $\frac{1}{x}$ of the tank per hour, Pipe B fills $\frac{1}{x+1}$ per hour. Working together, what fraction of the tank do they fill in one hour? Give the combined rate as a single simplified rational expression. _____

22. A trip of D miles is driven at speed r mph, then the same distance back at speed $r + 10$ mph. Express the total time, in hours, as a single simplified rational expression in r and D . _____

23. A baker uses $\frac{x}{x+2}$ cups of flour for one recipe and $\frac{3}{x+2}$ cups for another. Find the total flour used, in simplest form, and state the physical restriction on x . _____

24. In a parallel-resistor circuit, the total resistance satisfies $\frac{1}{R_{\text{total}}} = \frac{1}{R_1} + \frac{1}{R_2}$. With $R_1 = x$ ohms and $R_2 = x + 4$ ohms, find a single simplified expression for R_{total} . _____

Additional Practice

25. Simplify $\frac{x^2-9}{x-3}$. _____

26. Excluded value of $\frac{1}{x+4}$. _____

27. Domain of $f(x) = \frac{x}{x-5}$. _____

28. Multiply $\frac{x}{3} \cdot \frac{6}{x}$. _____

29. Divide $\frac{x^2}{5} \div \frac{x}{10}$. _____

30. Add $\frac{3}{x} + \frac{5}{x}$. _____



Scan Me

31. Subtract $\frac{7}{x-1} - \frac{2}{x-1}$. _____

32. Solve $\frac{1}{x} = 4$. _____

33. Solve $\frac{x+2}{x-1} = 3$. _____

34. Vertical asymptote of $y = \frac{4}{x+8}$. _____



Answer Keys

<p>1. $\frac{5}{x}, x \neq 0$</p> <p>2. $\frac{3}{x-1}, x \neq 1$</p> <p>3. $\frac{3x+1}{x(x+1)}, x \neq 0, -1$</p> <p>4. $\frac{x-7}{(x+2)(x-1)}, x \neq -2, 1$</p> <p>5. $(x-3)(x+3)$</p> <p>6. $\frac{x^2-2x-1}{x(x+1)(x-1)}, x \neq 0, \pm 1$</p> <p>7. $\frac{x^2-x+6}{(x-3)(x+3)}, x \neq \pm 3$</p> <p>8. $\frac{3x+14}{(x-4)(x+4)}, x \neq \pm 4$</p> <p>9. $\frac{x^2-4x+10}{(x-2)(x+1)}, x \neq 2, -1$</p> <p>10. $\frac{2x-1}{x(x+1)}, x \neq 0, -1$</p> <p>11. $\frac{3x+2}{(x-1)(x+1)}, x \neq \pm 1$</p> <p>12. $\frac{3x+1}{x-1}, x \neq 1$</p> <p>Additional Practice Answers</p> <p>25. $x+3, x \neq 3$</p> <p>26. $x = -4$</p> <p>27. $x \neq 5$</p> <p>28. 2</p> <p>29. $2x$</p>	<p>13. $\frac{2x}{(x-2)(x+2)}, x \neq \pm 2$</p> <p>14. $\frac{10x+5}{(x-2)(x+3)}, x \neq 2, -3$</p> <p>15. $\frac{7}{2x}, x \neq 0$</p> <p>16. FALSE</p> <p>17. $\frac{x+1}{x+3}, x \neq -3$</p> <p>18. $\frac{2x+2}{(x-2)(x+2)}, x \neq \pm 2$</p> <p>19. $\frac{1}{x-3}, x \neq 3$</p> <p>20. $\frac{x^2+3x+1}{x(x+1)}, x \neq 0, -1$</p> <p>21. $\frac{2x+1}{x(x+1)}$ per hour, $x > 0$</p> <p>22. $\frac{D(2r+10)}{r(r+10)}$ hours, $r > 0$</p> <p>23. $\frac{x+3}{x+2}$ cups, $x \geq 0$</p> <p>24. $R_{\text{total}} = \frac{x(x+4)}{2x+4}$ ohms, $x > 0$</p> <p>30. $\frac{8}{x}$</p> <p>31. $\frac{5}{x-1}$</p> <p>32. $x = \frac{1}{4}$</p> <p>33. $x = \frac{5}{2}$</p> <p>34. $x = -8$</p>
---	---

Additional Practice: Answers for all numbered items, including the added practice, are shown in the grid above.

Step-by-Step Explanations

- A careful way to see it: Same denominator: $\frac{2+3}{x} = \frac{5}{x}$. (The table values match $\frac{5}{x}$ at $x = 1, 2, 5$.) That gives a quick check on the answer.
- Same denominator: combine numerators. $(x+5)-(x+2) = x+5-x-2 = 3$. Watch the sign: it hits both x and -2 .
- LCD = $x(x+1)$. $\frac{x+1}{x(x+1)} + \frac{2x}{x(x+1)} = \frac{3x+1}{x(x+1)}$. (The table matches $\frac{3x+1}{x(x+1)}$ at $x = 1, 2, 3$.)
- LCD = $(x+2)(x-1)$. Numerators: $3(x-1)-2(x+2) = 3x-3-2x-4 = x-7$. (The table agrees with $\frac{x-7}{(x+2)(x-1)}$ at $x = 0, 2, 3$.)
- Factor: $x^2-9 = (x-3)(x+3)$. The LCD is $(x-3)(x+3)$ (each factor at its highest power).
- LCD = $x(x+1)(x-1)$. Numerators: $(x+1)(x-1)+x(x-1)-x(x+1) = (x^2-1)+(x^2-x)-(x^2+x) = x^2-2x-1$.
- One steady path is: LCD = $(x-3)(x+3)$. Numerators: $2(x+3)+x(x-3) = 2x+6+x^2-3x = x^2-x+6$. That gives a quick check on the answer.
- Start with the key idea: $x^2-16 = (x-4)(x+4)$. LCD = $(x-4)(x+4)$. $\frac{3(x+4)+2}{(x-4)(x+4)} = \frac{3x+14}{(x-4)(x+4)}$. That gives a quick check on the answer.
- A careful way to see it: LCD = $(x-2)(x+1)$. Numerators: $x(x+1)-5(x-$

- $2) = x^2+x-5x+10 = x^2-4x+10$. That gives a quick check on the answer.
- Keep the rule visible: LCD = $x(x+1)$. Numerators: $3x-(x+1) = 3x-x-1 = 2x-1$. This is the part to check before moving on, because it keeps the answer tied to the original question.
- One steady path is: $x^2-1 = (x-1)(x+1)$. LCD = $(x-1)(x+1)$. Numerators: $5+3(x-1) = 5+3x-3 = 3x+2$. Final: $\frac{3x+2}{(x-1)(x+1)}$. That gives a quick check on the answer.
- The denominators already match, so just add the tops: $2x+(x+1) = 3x+1$, over the shared $x-1$. The numerator doesn't factor to cancel anything, so keep $x \neq 1$.
- The LCD is $(x-2)(x+2)$. Rewrite each piece: $\frac{x-2}{(x-2)(x+2)} + \frac{x+2}{(x-2)(x+2)}$. Add the tops: $(x-2)+(x+2) = 2x$, giving $\frac{2x}{(x-2)(x+2)}$. Keep $x \neq \pm 2$.
- LCD = $(x-2)(x+3)$. Numerators: $(x+3)^2-(x-2)^2 = (x^2+6x+9)-(x^2-4x+4) = 10x+5$.
- One steady path is: LCD = $2x \cdot \frac{8-1}{2x} = \frac{7}{2x}$. This is the part to check before moving on, because it keeps the answer tied to the original question.
- Add fractions over a common denominator: $\frac{ad+bc}{bd}$. Adding numerators and denominators separately is one of the most universal fraction errors.



17. Denominators match, so subtract the tops – but the minus sign hits *both* terms of $(x - 1)$: $2x - (x - 1) = 2x - x + 1 = x + 1$. Result: $\frac{x+1}{x+3}$, with $x \neq -3$.

18. Keep the rule visible: $x^2 - 4 = (x - 2)(x + 2)$. LCD = $(x - 2)(x + 2)$. $\frac{x + (x + 2)}{(x - 2)(x + 2)} = \frac{2x + 2}{(x - 2)(x + 2)}$. That gives a quick check on the answer.

19. One steady path is: $3 - x = -(x - 3)$. So $\frac{4}{3-x} = -\frac{4}{x-3}$. Sum: $\frac{5-4}{x-3} = \frac{1}{x-3}$. (Sign-flip denominators are the same denominator in disguise.) That gives a quick check on the answer.

20. Start with the key idea: LCD = $x(x + 1)$. $\frac{(x + 1)^2 + x}{x(x + 1)} = \frac{x^2 + 2x + 1 + x}{x(x + 1)} = \frac{x^2 + 3x + 1}{x(x + 1)}$. That gives a quick check on the answer.

21. Combined rate = $\frac{1}{x} + \frac{1}{x+1}$. LCD = $x(x + 1)$. $\frac{(x + 1) + x}{x(x + 1)} = \frac{2x + 1}{x(x + 1)}$ per hour. The algebraic restriction is $x \neq 0, -1$; physically the rates must be positive, so $x > 0$. Sanity check at $x = 2$: $\frac{1}{2} + \frac{1}{3} = \frac{5}{6}$. Simplified: $\frac{5}{6}$. Match. (At $x = 2$ the two pipes together fill $\frac{5}{6}$ of the tank per hour – so they finish a full

tank in $\frac{6}{5} = 1.2$ hours.)

22. Time = distance/speed. Out: $\frac{D}{r}$. Back: $\frac{D}{r+10}$. Total: $\frac{D}{r} + \frac{D}{r+10} = \frac{D(r+10) + Dr}{r(r+10)} = \frac{D(2r+10)}{r(r+10)}$. Restriction: $r \neq 0, -10$; physically $r > 0$ (and $r + 10 > 0$ automatically). Sanity check with $D = 60$, $r = 30$:

$\frac{60}{30} + \frac{60}{40} = 2 + 1.5 = 3.5$ h. Simplified: $\frac{60 \cdot 70}{30 \cdot 40} = \frac{4200}{1200} = 3.5$ h. Match.

23. Same denominator: add the numerators. $\frac{x+3}{x+2}$ cups total. Algebraic restriction: $x \neq -2$; physical restriction (cups can't be negative): $x \geq 0$. Sanity check at $x = 4$: $\frac{4}{6} + \frac{3}{6} = \frac{7}{6}$ cups. Simplified: $\frac{7}{6}$. Match.

24. Start with the key idea: $\frac{1}{R_{\text{total}}} = \frac{1}{x} + \frac{1}{x+4} = \frac{(x+4) + x}{x(x+4)} = \frac{2x+4}{x(x+4)}$.

Take the reciprocal: $R_{\text{total}} = \frac{x(x+4)}{2x+4}$ ohms. Algebraic restrictions: $x \neq 0, -4$;

physically $x > 0$. Sanity check at $x = 4$: $\frac{1}{4} + \frac{1}{8} = \frac{3}{8}$, so $R = \frac{8}{3} \approx 2.67$ ohms.

Formula: $\frac{4 \cdot 8}{12} = \frac{32}{12} = \frac{8}{3}$. Match. (Parallel resistance is always less than either individual resistor – here $\frac{8}{3} < 4$ and $\frac{8}{3} < 8$.) That gives a quick check on the answer.



Build Algebra Confidence From Pre-Algebra Through Algebra II



The Complete Algebra Success Bundle

Pre-Algebra, Algebra I, and Algebra II in one clear path

Friendly lessons, focused practice, and full-review support for every stage.



Scan for the Bundle

6 Books
3 Courses
1 Path

Bundle Value: Six coordinated books help students review missing skills, learn new algebra topics, and practice until the steps feel natural.

Complete Course Path

- ✓ Starts with Pre-Algebra foundations
- ✓ Moves smoothly into Algebra I skills
- ✓ Extends learning through Algebra II topics
- ✓ Great for review, tutoring, and summer study

One bundle, one steady path.

Step-by-Step Lessons

- ✓ Plain-English explanations students can follow
- ✓ Worked examples that show every important step
- ✓ Common mistakes called out before they stick
- ✓ Skill-building practice after each lesson
- ✓ Helpful for independent study or class support

Less guessing. More understanding.

Practice That Sticks

- ✓ Matching practice workbooks for extra repetition
- ✓ Review sets to keep older skills fresh
- ✓ Answer explanations for checking thinking
- ✓ Strong support before tests and final exams
- ✓ Designed to build fluency and confidence

Practice today. Remember tomorrow.

STUDENT FAVORITE • Master Algebra II From the Ground Up



Algebra II for Beginners

Written by a top math teacher & aligned with national and state Algebra II courses. From polynomial functions to logarithms, trigonometry, and rational expressions — explained the easy way.

- ✓ **Complete coverage** of every Algebra II concept — perfect companion to these worksheets
- ✓ **Step-by-step explanations** with worked examples on every topic
- ✓ **QR codes in every chapter** for free video lessons & bonus practice
- ✓ **2 full-length practice tests** with detailed answer keys

- ✓ 100% Guaranteed
- ✓ Lifetime Support
- ✓ Trusted by Teachers

Start Your Algebra Journey Today! →

★ STUDENT'S #1 CHOICE ★

Teacher-recommended • 12,000+ Happy Students

PDF EDITION



Instant download • any device

PAPERBACK



Paperback on Amazon

Hold it in your hands

Pair these free worksheets with *Algebra II for Beginners* and you have a complete self-paced course — concept lessons, daily practice, and full exam-style reviews, all in one path. → EffortlessMath.com/product/algebra-ii-for-beginners