

Weighted Averages and Missing Data Values

Name: _____ Date: _____ Score: _____ / 18

Quick Review and Helpful Hints

A *weighted average* multiplies each value by its weight (or count), adds the products, then divides by the total weight. To find a *missing value* when the mean is known, multiply the mean by the number of values to get the total, then subtract the values you know.

▶ **Example:** A student scores 80 on 2 tests and 95 on 3 tests. Find the weighted average. **Work:** $\frac{80(2) + 95(3)}{2 + 3} = \frac{160 + 285}{5} = \frac{445}{5}$.

$$\frac{\sum(\text{value} \times \text{weight})}{\sum \text{weight}}$$

★ **Answer:** 89

Multiply, add, then divide.

Practice Problems

Find each average or missing value.

- | | |
|---|---|
| 1. Mean of 4, 6, 8
_____ | 8. Mean of 2, 4, 6, 8, 10
_____ |
| 2. Mean of 10, 20, 30, 40
_____ | 9. Mean of 4 values is 10; three are 8, 10, 12. Find the missing value
_____ |
| 3. $\frac{90(1) + 80(1)}{2}$
_____ | 10. $\frac{100(3) + 50(1)}{4}$
_____ |
| 4. $\frac{70(2) + 100(3)}{5}$
_____ | 11. Mean of 12, 18
_____ |
| 5. Mean of 3 tests is 80; two are 75, 85. Find the third
_____ | 12. Mean of two is 20; one is 15. Find the other
_____ |
| 6. Mean of 5, 5, 5, 5
_____ | 13. $\frac{80(2) + 90(2)}{4}$
_____ |
| 7. $\frac{60(1) + 90(2)}{3}$
_____ | 14. Mean of 3, 7, 11
_____ |

Word Problems

15. Grades: 85 on 2 tests and 95 on 2 tests. Find the weighted average.

16. A class of 10 averages 70; a class of 20 averages 85. Find the combined average.

17. Five quiz scores average 8; four of them are 7, 9, 8, 10. Find the fifth.

18. Homework counts 40% and the exam 60%. Homework = 90, exam = 80. Find the final grade.



Answer Keys

- | | | |
|------------------------------------|---------------------------------------|-------------------------------------|
| 1. <input type="text" value="6"/> | 7. <input type="text" value="80"/> | 13. <input type="text" value="85"/> |
| 2. <input type="text" value="25"/> | 8. <input type="text" value="6"/> | 14. <input type="text" value="7"/> |
| 3. <input type="text" value="85"/> | 9. <input type="text" value="10"/> | 15. <input type="text" value="90"/> |
| 4. <input type="text" value="88"/> | 10. <input type="text" value="87.5"/> | 16. <input type="text" value="80"/> |
| 5. <input type="text" value="80"/> | 11. <input type="text" value="15"/> | 17. <input type="text" value="6"/> |
| 6. <input type="text" value="5"/> | 12. <input type="text" value="25"/> | 18. <input type="text" value="84"/> |

Step-by-Step Explanations

1. Start by naming the process: For an average, add the values and divide by how many values there are; for a missing value, work backward from the total. The setup/work is $\frac{4 + 6 + 8}{3} = \frac{18}{3} = 6$. So the final answer is 6.
2. A good way to think about this is: For an average, add the values and divide by how many values there are; for a missing value, work backward from the total. The setup/work is $\frac{100}{4} = 25$. So the final answer is 25.
3. Step by step: For an average, add the values and divide by how many values there are; for a missing value, work backward from the total. The setup/work is $\frac{90 + 80}{2} = 85$. So the final answer is 85.
4. Take it one move at a time: For an average, add the values and divide by how many values there are; for a missing value, work backward from the total. The setup/work is $\frac{140 + 300}{5} = \frac{440}{5} = 88$. So the final answer is 88.
5. Start by naming the process: For an average, add the values and divide by how many values there are; for a missing value, work backward from the total. The setup/work is Total = $80 \times 3 = 240$. The two known scores sum to 160, so the third = $240 - 160 = 80$. So the final answer is 80.
6. A good way to think about this is: For an average, add the values and divide by how many values there are; for a missing value, work backward from the total. The setup/work is All values are 5: mean 5. So the final answer is 5.
7. Step by step: For an average, add the values and divide by how many values there are; for a missing value, work backward from the total. The setup/work is $\frac{60 + 180}{3} = \frac{240}{3} = 80$. So the final answer is 80.
8. Take it one move at a time: For an average, add the values and divide by how many values there are; for a missing value, work backward from the total. The setup/work is $\frac{30}{5} = 6$. So the final answer is 6.
9. Start by naming the process: For an average, add the values and divide by how many values there are; for a missing value, work backward from the total. The setup/work is Total = $10 \times 4 = 40$. Subtract 30: missing = 10. So the final answer is 10.
10. A good way to think about this is: For an average, add the values and divide by how many values there are; for a missing value, work backward from the total. The setup/work is $\frac{300 + 50}{4} = \frac{350}{4} = 87.5$. So the final answer is 87.5.
11. Step by step: For an average, add the values and divide by how many values there are; for a missing value, work backward from the total. The setup/work is $\frac{12 + 18}{2} = 15$. So the final answer is 15.
12. Take it one move at a time: For an average, add the values and divide by how many values there are; for a missing value, work backward from the total. The setup/work is Total = $20 \times 2 = 40$. Subtract 15: other = 25. So the final answer is 25.
13. Start by naming the process: For an average, add the values and divide by how many values there are; for a missing value, work backward from the total. The setup/work is $\frac{160 + 180}{4} = \frac{340}{4} = 85$. So the final answer is 85.
14. A good way to think about this is: For an average, add the values and divide by how many values there are; for a missing value, work backward from the total. The setup/work is $\frac{3 + 7 + 11}{3} = \frac{21}{3} = 7$. So the final answer is 7.
15. Step by step: For an average, add the values and divide by how many values there are; for a missing value, work backward from the total. The setup/work is $\frac{85(2) + 95(2)}{4} = \frac{170 + 190}{4} = 90$. So the final answer is 90.
16. Take it one move at a time: For an average, add the values and divide by how many values there are; for a missing value, work backward from the total. The setup/work is $\frac{10(70) + 20(85)}{30} = \frac{2400}{30} = 80$. So the final answer is 80.
17. Start by naming the process: For an average, add the values and divide by how many values there are; for a missing value, work backward from the total. The setup/work is Total = $8 \times 5 = 40$. Subtract 34: fifth = 6. So the final answer is 6.
18. A good way to think about this is: For an average, add the values and divide by how many values there are; for a missing value, work backward from the total. The setup/work is $0.4(90) + 0.6(80) = 36 + 48 = 84$. So the final answer is 84.



Keep Building CBEST Math Skills

Recommended Effortless Math resources



CBEST Math Test Prep Bundle

Use the complete CBEST Math resource for review, worked examples, extra practice, and test-style questions after each worksheet.



Scan Me
Download Instantly

STUDENT FAVORITE - CBEST Math for Beginners



CBEST Math for Beginners 2026

Step-by-step lessons, topic practice, and full review support for students who want a calm path through CBEST Math preparation.

A strong companion for self-study, tutoring, homework, and targeted review.

PDF Edition



Scan Me
Download Instantly

For more CBEST Math prep, visit EffortlessMath.com/CBEST