


Literal Equations and Formulas

Name: _____ Date: _____ Score: _____ / 18

Quick Review and Helpful Hints

A *literal equation* contains more than one variable (often a formula). To solve for one variable, isolate it with inverse operations, treating the other letters as if they were numbers. The result lets you plug in values any time.

▶ **Example:** Solve $A = \frac{1}{2}bh$ for h . **Work:** Multiply both sides by 2:
 $2A = bh$. Divide both sides by b : $h = \frac{2A}{b}$. ★ **Answer:** $h = \frac{2A}{b}$



$A = \frac{1}{2}bh$, so $h = \frac{2A}{b}$.

Practice Problems

Solve each for the indicated variable.

- | | | | |
|--------------------------|-------|--------------------------------|-------|
| 1. $d = rt$ for r | _____ | 8. $A = \frac{1}{2}bh$ for b | _____ |
| 2. $A = lw$ for w | _____ | 9. $V = lwh$ for h | _____ |
| 3. $P = 4s$ for s | _____ | 10. $F = ma$ for a | _____ |
| 4. $C = 2\pi r$ for r | _____ | 11. $I = Prt$ for t | _____ |
| 5. $y = mx + b$ for b | _____ | 12. $2x + y = 10$ for y | _____ |
| 6. $y = mx + b$ for m | _____ | 13. $ax = b$ for x | _____ |
| 7. $P = 2l + 2w$ for l | _____ | 14. $\frac{x}{c} = d$ for x | _____ |

Word Problems

15. The area of a rectangle is $A = lw$. Solve for l , then find l when $A = 24$ and $w = 4$. _____
16. Distance is $d = rt$. Solve for t , then find t when $d = 120$ and $r = 40$. _____
17. Perimeter is $P = 2l + 2w$. Solve for w , then find w when $P = 20$ and $l = 6$. _____
18. Simple interest is $I = Prt$. Solve for r , then find r when $I = 60$, $P = 500$, $t = 2$. _____



Answer Keys

1. $r = \frac{d}{t}$

2. $w = \frac{A}{l}$

3. $s = \frac{P}{4}$

4. $r = \frac{C}{2\pi}$

5. $b = y - mx$

6. $m = \frac{y - b}{x}$

7. $l = \frac{P - 2w}{2}$

8. $b = \frac{2A}{h}$

9. $h = \frac{V}{lw}$

10. $a = \frac{F}{m}$

11. $t = \frac{I}{Pr}$

12. $y = 10 - 2x$

13. $x = \frac{b}{a}$

14. $x = cd$

15. $l = \frac{A}{w} = 6$

16. $t = \frac{d}{r} = 3$

17. $w = \frac{P - 2l}{2} = 4$

18. $r = \frac{I}{Pt} = 0.06$

Step-by-Step Explanations

1. Start by naming the process: Read what the problem is asking, choose the matching rule, write the setup, and then simplify one step at a time. The setup/work is Divide both sides by t : $r = \frac{d}{t}$. So the final answer is $r = \frac{d}{t}$.

2. A good way to think about this is: Read what the problem is asking, choose the matching rule, write the setup, and then simplify one step at a time. The setup/work is Divide both sides by l : $w = \frac{A}{l}$. So the final answer is $w = \frac{A}{l}$.

3. Step by step: Read what the problem is asking, choose the matching rule, write the setup, and then simplify one step at a time. The setup/work is Divide both sides by 4: $s = \frac{P}{4}$. So the final answer is $s = \frac{P}{4}$.

4. Take it one move at a time: Read what the problem is asking, choose the matching rule, write the setup, and then simplify one step at a time. The setup/work is Divide both sides by 2π : $r = \frac{C}{2\pi}$. So the final answer is $r = \frac{C}{2\pi}$.

5. Start by naming the process: Read what the problem is asking, choose the matching rule, write the setup, and then simplify one step at a time. The setup/work is Subtract mx from both sides: $b = y - mx$. So the final answer is $b = y - mx$.

6. A good way to think about this is: Read what the problem is asking, choose the matching rule, write the setup, and then simplify one step at a time. The setup/work is Subtract b : $y - b = mx$. Divide by x : $m = \frac{y - b}{x}$. So the final answer is $m = \frac{y - b}{x}$.

7. Step by step: Read what the problem is asking, choose the matching rule, write the setup, and then simplify one step at a time. The setup/work is Subtract $2w$: $P - 2w = 2l$. Divide by 2: $l = \frac{P - 2w}{2}$. So the final answer is $l = \frac{P - 2w}{2}$.

8. Take it one move at a time: Read what the problem is asking, choose the matching rule, write the setup, and then simplify one step at a time. The setup/work is Multiply by 2: $2A = bh$. Divide by h : $b = \frac{2A}{h}$. So the final answer is $b = \frac{2A}{h}$.

9. Start by naming the process: Read what the problem is asking, choose the matching rule, write the setup, and then simplify one step at a time. The setup/work is Divide both sides by lw : $h = \frac{V}{lw}$. So the final answer is $h = \frac{V}{lw}$.

10. A good way to think about this is: Read what the problem is asking, choose the matching rule, write the setup, and then simplify one step at a time. The setup/work is Divide both sides by m : $a = \frac{F}{m}$. So the final answer is $a = \frac{F}{m}$.

11. Step by step: Read what the problem is asking, choose the matching rule, write the setup, and then simplify one step at a time. The setup/work is Divide both sides by Pr : $t = \frac{I}{Pr}$. So the final answer is $t = \frac{I}{Pr}$.

12. Take it one move at a time: Read what the problem is asking, choose the matching rule, write the setup, and then simplify one step at a time. The setup/work is Subtract $2x$ from both sides: $y = 10 - 2x$. So the final answer is $y = 10 - 2x$.

13. Start by naming the process: Read what the problem is asking, choose the matching rule, write the setup, and then simplify one step at a time. The setup/work is Divide both sides by a : $x = \frac{b}{a}$. So the final answer is $x = \frac{b}{a}$.

14. A good way to think about this is: Read what the problem is asking, choose the matching rule, write the setup, and then simplify one step at a time. The setup/work is Multiply both sides by c : $x = cd$. So the final answer is $x = cd$.

15. Step by step: Read what the problem is asking, choose the matching rule, write the setup, and then simplify one step at a time. The setup/work is $l = \frac{A}{w} = \frac{24}{4} = 6$. So the final answer is $l = \frac{A}{w} = 6$.

16. Take it one move at a time: Read what the problem is asking, choose the matching rule, write the setup, and then simplify one step at a time. The setup/work is $t = \frac{d}{r} = \frac{120}{40} = 3$. So the final answer is $t = \frac{d}{r} = 3$.

17. Start by naming the process: Read what the problem is asking, choose the matching rule, write the setup, and then simplify one step at a time. The setup/work is $w = \frac{P - 2l}{2} = \frac{20 - 12}{2} = 4$. So the final answer is $w = \frac{P - 2l}{2} = 4$.

18. A good way to think about this is: Read what the problem is asking, choose the matching rule, write the setup, and then simplify one step at a time. The setup/work is $r = \frac{I}{Pt} = \frac{60}{1000} = 0.06$. So the final answer is $r = \frac{I}{Pt} = 0.06$.



Keep Building Praxis Core Math Skills

Recommended Effortless Math resources




The Most Comprehensive Praxis Core Math Preparation Bundle

Use the complete Praxis Core Math resource for review, worked examples, extra practice, and test-style questions after each worksheet.



Scan Me
Download Instantly

STUDENT FAVORITE - Praxis Core Math for Beginners



Praxis Core Math for Beginners 2026

Step-by-step lessons, topic practice, and full review support for students who want a calm path through Praxis Core Math preparation.

A strong companion for self-study, tutoring, homework, and targeted review.

PDF Edition



Scan Me
Download Instantly

For more Praxis Core Math prep, visit EffortlessMath.com/Praxis-Core