

The Quadratic Formula

Name: _____ Date: _____ Score: _____ / 18

Quick Review and Helpful Hints

For $ax^2 + bx + c = 0$, the solutions are $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$. Identify a, b, c , then substitute carefully. The *discriminant* $b^2 - 4ac$ tells how many real solutions: positive gives two, zero gives one, negative gives none.

▶ **Example:** Solve $x^2 + 5x + 6 = 0$ with the formula. **Work:** $a = 1, b = 5, c = 6$. Discriminant $= 25 - 24 = 1$. $x = \frac{-5 \pm 1}{2}$. ★ **Answer:** $x = -2$ or $x = -3$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Identify a, b, c , then substitute.

◆ **Practice Problems**

Solve, or find the discriminant, as directed.

- | | |
|---|--|
| <p>1. $x^2 + 5x + 6 = 0$ _____</p> <p>2. $x^2 - 3x - 4 = 0$ _____</p> <p>3. $x^2 + x - 12 = 0$ _____</p> <p>4. $x^2 - 5x + 6 = 0$ _____</p> <p>5. $x^2 - 2x - 3 = 0$ _____</p> <p>6. $x^2 + 7x + 12 = 0$ _____</p> <p>7. $x^2 - 9 = 0$ _____</p> | <p>8. $x^2 - 6x + 5 = 0$ _____</p> <p>9. $x^2 + 2x - 15 = 0$ _____</p> <p>10. $x^2 - 7x + 10 = 0$ _____</p> <p>11. $x^2 + 4x + 3 = 0$ _____</p> <p>12. $x^2 - x - 20 = 0$ _____</p> <p>13. Discriminant of $x^2 + 3x + 2$ _____</p> <p>14. Discriminant of $x^2 - 4x + 4$ _____</p> |
|---|--|

◆ **Word Problems**

15. A quadratic revenue model is set equal to zero: $x^2 + 6x + 8 = 0$. Use the formula to find the break-even x -values. _____
16. Before graphing $x^2 + 5x + 6$, a student checks the discriminant. Find it and use it to decide how many real solutions there are. _____
17. A walkway model leads to $x^2 - 8x + 12 = 0$. Use the quadratic formula to solve for x . _____
18. How many real solutions are there if the discriminant equals 0? _____



Answer Keys

1. $x = -2, -3$

2. $x = 4, -1$

3. $x = 3, -4$

4. $x = 2, 3$

5. $x = 3, -1$

6. $x = -3, -4$

7. $x = 3, -3$

8. $x = 1, 5$

9. $x = 3, -5$

10. $x = 2, 5$

11. $x = -1, -3$

12. $x = 5, -4$

13. 1

14. 0

15. $x = -2, -4$

16. 1

17. $x = 2, 6$

18. one

Step-by-Step Explanations

1. Start by naming the process: Use the quadratic formula step by step: identify a , b , and c , compute the discriminant, then simplify. The setup/work is $b^2 - 4ac = 25 - 24 = 1$; $x = \frac{-5 \pm 1}{2} = -2, -3$. So the final answer is $x = -2, -3$.

2. A good way to think about this is: Use the quadratic formula step by step: identify a , b , and c , compute the discriminant, then simplify. The setup/work is $9 + 16 = 25$; $x = \frac{3 \pm 5}{2} = 4, -1$. So the final answer is $x = 4, -1$.

3. Step by step: Use the quadratic formula step by step: identify a , b , and c , compute the discriminant, then simplify. The setup/work is $1 + 48 = 49$; $x = \frac{-1 \pm 7}{2} = 3, -4$. So the final answer is $x = 3, -4$.

4. Take it one move at a time: Use the quadratic formula step by step: identify a , b , and c , compute the discriminant, then simplify. The setup/work is $25 - 24 = 1$; $x = \frac{5 \pm 1}{2} = 3, 2$. So the final answer is $x = 2, 3$.

5. Start by naming the process: Use the quadratic formula step by step: identify a , b , and c , compute the discriminant, then simplify. The setup/work is $4 + 12 = 16$; $x = \frac{2 \pm 4}{2} = 3, -1$. So the final answer is $x = 3, -1$.

6. A good way to think about this is: Use the quadratic formula step by step: identify a , b , and c , compute the discriminant, then simplify. The setup/work is $49 - 48 = 1$; $x = \frac{-7 \pm 1}{2} = -3, -4$. So the final answer is $x = -3, -4$.

7. Step by step: Use the quadratic formula step by step: identify a , b , and c , compute the discriminant, then simplify. The setup/work is $a = 1, b = 0, c = -9$; $x = \frac{\pm 6}{2} = 3, -3$. So the final answer is $x = 3, -3$.

8. Take it one move at a time: Use the quadratic formula step by step: identify a , b , and c , compute the discriminant, then simplify. The setup/work is $36 - 20 = 16$; $x = \frac{6 \pm 4}{2} = 5, 1$. So the final answer is $x = 1, 5$.

9. Start by naming the process: Use the quadratic formula step by step: identify a , b , and c , compute the discriminant, then simplify. The setup/work is $4 + 60 = 64$; $x = \frac{-2 \pm 8}{2} = 3, -5$. So the final answer is $x = 3, -5$.

10. A good way to think about this is: Use the quadratic formula step by step: identify a , b , and c , compute the discriminant, then simplify. The setup/work is $49 - 40 = 9$; $x = \frac{7 \pm 3}{2} = 5, 2$. So the final answer is $x = 2, 5$.

11. Step by step: Use the quadratic formula step by step: identify a , b , and c , compute the discriminant, then simplify. The setup/work is $16 - 12 = 4$; $x = \frac{-4 \pm 2}{2} = -1, -3$. So the final answer is $x = -1, -3$.

12. Take it one move at a time: Use the quadratic formula step by step: identify a , b , and c , compute the discriminant, then simplify. The setup/work is $1 + 80 = 81$; $x = \frac{1 \pm 9}{2} = 5, -4$. So the final answer is $x = 5, -4$.

13. Start by naming the process: Use the quadratic formula step by step: identify a , b , and c , compute the discriminant, then simplify. The setup/work is $b^2 - 4ac = 9 - 8 = 1$. So the final answer is 1.

14. A good way to think about this is: Use the quadratic formula step by step: identify a , b , and c , compute the discriminant, then simplify. The setup/work is $b^2 - 4ac = 16 - 16 = 0$. So the final answer is 0.

15. Step by step: Use the quadratic formula step by step: identify a , b , and c , compute the discriminant, then simplify. The setup/work is $36 - 32 = 4$; $x = \frac{-6 \pm 2}{2} = -2, -4$. So the final answer is $x = -2, -4$.

16. Take it one move at a time: Use the quadratic formula step by step: identify a , b , and c , compute the discriminant, then simplify. The setup/work is $25 - 24 = 1$. So the final answer is 1.

17. Start by naming the process: Use the quadratic formula step by step: identify a , b , and c , compute the discriminant, then simplify. The setup/work is $64 - 48 = 16$; $x = \frac{8 \pm 4}{2} = 6, 2$. So the final answer is $x = 2, 6$.

18. A good way to think about this is: Use the quadratic formula step by step: identify a , b , and c , compute the discriminant, then simplify. The setup/work is A discriminant of 0 gives exactly one real solution. So the final answer is one.



Keep Building Praxis Core Math Skills

Recommended Effortless Math resources



The Most Comprehensive Praxis Core Math Preparation Bundle

Use the complete Praxis Core Math resource for review, worked examples, extra practice, and test-style questions after each worksheet.



Scan Me
Download Instantly

STUDENT FAVORITE - Praxis Core Math for Beginners



Praxis Core Math for Beginners 2026

Step-by-step lessons, topic practice, and full review support for students who want a calm path through Praxis Core Math preparation.

A strong companion for self-study, tutoring, homework, and targeted review.

PDF Edition



Scan Me
Download Instantly

For more Praxis Core Math prep, visit [EffortlessMath.com/Praxis-Core](https://www.EffortlessMath.com/Praxis-Core)